



## Classification for Fish Sampling<sup>1</sup>

This standard is issued under the fixed designation D 4211; the number immediately following the designation indicates the year of original adoption or, in the case of revision, the year of last revision. A number in parentheses indicates the year of last reapproval. A superscript epsilon (ε) indicates an editorial change since the last revision or reapproval.

### 1. Scope

1.1 This classification covers rotenone and antimycin which are used to collect or eradicate fish; numerous chemicals have been used but presently only rotenone and antimycin are EPA approved for this use.

### 2. Referenced Document

2.1 *ASTM Standards*:<sup>2</sup>

D 4131 Practice for Sampling Fish with Rotenone

### 3. Significance and Use

3.1 The significance of using chemical fish toxicants is that more complete population analyses or total eradication, or both, can be accomplished. Target species can be selectively eradicated by varying concentrations. This provides a very effective tool in fisheries investigations and management programs. Water conditions (that is, pH, temperature, alkalinity, and so forth) and morphology can be limiting factors.

3.2 *Rotenone*—Rotenone used as a fish toxicant is highly versatile and can be used effectively to collect fish samples; to eradicate fish; and to selectively remove certain fish species.

3.2.1 Its effectiveness is reduced in cold <20°C, and dosage required increases with alkalinities. It may also eliminate food web organisms. Fish may be repulsed from treated areas.

3.3 *Antimycin*—Antimycin is versatile in the selective removal of scalefish or even more selectively against certain centrarchids (sunfish) and minnows.

3.3.1 Its effectiveness is reduced in water with pH above 8.5.

### 4. Basis of Classification

#### 4.1 *EPA-Approved Fish Toxicants*.

4.1.1 *Rotenone*—This fish toxicant is also known as derris or cube and is derived from roots of several plants of the family *Leguminosae*. Its action mode as a powerful respiratory inhibitor in fish starts with entrance into the blood stream via the gills and then by translocation to vital organs. Formulations are in the following general forms: a liquid containing 5 % rotenone, liquid with 2.5 % rotenone and synergists, and wettable powder. Persistence in the environment is seldom more than two weeks although it may remain longer in very cold, soft water.

4.1.2 *Antimycin*—This fish-toxicant is an antibiotic produced by *Streptomyces* and is known as Fintrol 5®, Fintrol 15®, and Fintrol-Concentrate®. The Fintrol 5® and 15® are coated sand-like grains for use in waters 5 to 15 ft (1.52 to 4.57 m) deep. The Fintrol-Concentrate® is a liquid for use in streams or shallow waters. Antimycin is effective in small concentrations (5 ppb) and enters fish through the gills to irreversibly block cellular respiration. It is nonpersistent in the environment.

<sup>1</sup> This classification is under the jurisdiction of ASTM Committee E47 on Biological Effects and Environmental Fate and is the direct responsibility of Subcommittee E47.01 on Aquatic Assessment and Toxicology.

Current edition approved April 1, 2003. Published April 2003. Originally approved in 1982. Last previous edition approved in 1998 as D 4211 – 82 (1998).

<sup>2</sup> For referenced ASTM standards, visit the ASTM website, [www.astm.org](http://www.astm.org), or contact ASTM Customer Service at [service@astm.org](mailto:service@astm.org). For *Annual Book of ASTM Standards* volume information, refer to the standard's Document Summary page on the ASTM website.

*ASTM International takes no position respecting the validity of any patent rights asserted in connection with any item mentioned in this standard. Users of this standard are expressly advised that determination of the validity of any such patent rights, and the risk of infringement of such rights, are entirely their own responsibility.*

*This standard is subject to revision at any time by the responsible technical committee and must be reviewed every five years and if not revised, either reapproved or withdrawn. Your comments are invited either for revision of this standard or for additional standards and should be addressed to ASTM International Headquarters. Your comments will receive careful consideration at a meeting of the responsible technical committee, which you may attend. If you feel that your comments have not received a fair hearing you should make your views known to the ASTM Committee on Standards, at the address shown below.*

*This standard is copyrighted by ASTM International, 100 Barr Harbor Drive, PO Box C700, West Conshohocken, PA 19428-2959, United States. Individual reprints (single or multiple copies) of this standard may be obtained by contacting ASTM at the above address or at 610-832-9585 (phone), 610-832-9555 (fax), or [service@astm.org](mailto:service@astm.org) (e-mail); or through the ASTM website ([www.astm.org](http://www.astm.org)).*